. Appl. No.

: 09/748,706

Filed:

**December 22, 2000** 

## AMENDMENTS TO THE CLAIMS

Claims 1-39 (Cancelled)

- 40. (Currently Amended) A method of decoding the position of bioactive agents attached to microspheres on an array substrate comprising:
- a) providing an array <u>substrate composition</u> comprising a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation comprise:
  - i) a bioactive agent;
- ii) at least a first and a second decoding attribute wherein a first of said decoding attributes is an identifier binding ligand (IBL), wherein said first and second IBLs is are different from said bioactive agent and wherein said identifier binding ligand IBLs is are directly attached to said microsphere; and
- b) detecting each of said both the first and second decoding attributes IBLs to decode the position of identify each of said bioactive agents on said substrate.
- 41. (Currently Amended) The method according to claim 40, wherein said detecting comprises detecting binding of a <u>first</u> decoder binding ligand (DBL) to said <u>first</u> IBL <u>and a second decoder binding ligand to said second IBL</u>.
- 42. (Cancelled)
- 43. (Currently Amended) The method according to claims Claim 40 or 42, wherein said first and second IBLs comprise a nucleic acid.
- 44. (Cancelled)
- 45. (Currently Amended) The method according to elaims Claim 40 or 42, wherein said first and second attributes are IBLs and are attached to said first subpopulation of microspheres at a first ratio and are attached to said second population of microspheres at a second ratio.

Appl. No.

99/748,706

Filed:

December 22, 2000

45. (Currently Amended) The method according to claims Claim 40 or 42, wherein said first and second attributes are IBLs and are attached to said first subpopulation of microspheres at a first ratio and are attached to said second population of microspheres at a second ratio.

Claims 46-54 (Cancelled)

- 55. (New) A method of decoding the position of a bioactive agent on an array substrate comprising:
  - a) providing an array substrate comprising randomly distributed microspheres, wherein each microsphere comprises:
    - i. a bioactive agent; and
    - ii. a combination of different identifier binding ligands (IBLs), wherein said combination uniquely identifies said bioactive agent;
    - b) adding a first decoder binding ligand (DBL) to the array substrate, wherein the first DBL binds to an IBL in said combination;
    - c) identifying the position of the IBL that binds to the first DBL;
    - d) adding a second DBL to the array, wherein the second DBL binds to another IBL in said combination;
    - e) identifying the position of the IBL that binds to the second DBL; and
    - decoding the position of said bioactive agent on said array substrate based on the position on the array substrate of the IBLs that bind the first and the second DBLs.
- 56. (New) The method of Claim 55, wherein said combination of IBLs and said first and second DBLs comprise single strand nucleic acid.
- 57. (New) The method of Claim 56, wherein said nucleic acid is DNA.
- 58. (New) The method of Claim 56, wherein said nucleic acid is RNA.

Appl. No.

09/748,706

Filed:

**December 22, 2000** 

59. (New) The method of Claim 56, wherein said combination of IBLs and said first and second DBLs are oligonucleotides of about 8 to about 40 basepairs in length.

- 60. (New) The method of Claim 55, wherein said microspheres comprise 10 IBLs.
- 61. (New) The method of Claim 55, wherein said first DBL binds to said IBL with a dissociation constant of less than about 10<sup>-5</sup>-10<sup>-9</sup> M<sup>-1</sup>.
- 62. (New) The method of Claim 55, wherein said second DBL binds to said IBL with a dissociation constant of less than about 10<sup>-5</sup>-10<sup>-9</sup> M<sup>-1</sup>.
- 63. (New) The method of Claim 55, wherein said first and second DBLs comprise a fluorophore.